

Application No.: 10/028,416

AMENDMENTS TO THE CLAIMS:

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (currently amended) A method for predicting a polyadenylation site comprising: inputting a plurality of RNA transcript sequences ~~or sequences derived from RNA transcript sequences~~, wherein at least one sequence has its poly A or poly T tract sequence; searching for a polyadenylation site, wherein the polyadenylation is an adenine rich region at the 3' end of the sequence or a thymine rich region at the ~~beginning~~ 5' end of the sequence; detecting the presence of polyadenylation signals neighboring the polyadenylation site by ~~scanning~~ analyzing ~~[[the]]~~ EST or RNA sequences or their corresponding genomic DNA sequences.
2. (currently amended) The method of Claim 1 wherein the step of searching for a polyadenylation site ~~comprising~~ comprises scanning analyzing the sequences for adenine rich region at the 3' end of the sequence or a thymine rich region at the ~~beginning~~ 5' end of the sequence.

BEST AVAILABLE COPY

Application No.: 10/028,416

3. (original) The method of Claim 2 wherein the adenine rich region comprises adenine in at least 50% of the region and the thymine rich region comprises thymine in at least 50% of the region.
4. (original) The method of Claim 2 wherein the adenine rich region comprises adenine in at least 60% of the region and the thymine rich region comprises thymine in at least 60% of the region.
5. (original) The method of Claim 2 wherein the adenine rich region comprises adenine in at least 70% of the region and the thymine rich region comprises thymine in at least 70% of the region.
6. (original) The method of Claim 2 wherein the adenine rich region comprises adenine in at least 80% of the region and the thymine rich region comprises thymine in at least 80% of the region.
7. (currently amended) The method of Claim 1 wherein a heuristic score $n_A / (n_A + 0.5 * (\max(n_R - 20, 0)))$ is used for detecting adenine rich or thymine rich region; wherein n_A is the number of adenines or thymine in the block, and n_R is the number of bases after downstream of the block of adenines or thymine ~~thymine~~ to the end of the sequence.

Application No.: 10/028,416

8. (currently amended) A method for detecting polyadenylation signal in a sequence with a polyadenylation site comprising searching for a polyadenylation signal hexamer in the sequence before 5' of the polyadenylation site.

9. (currently amended) The method of Claim 8 wherein the searching comprises evaluating the probability that there is a polyadenylation [[site]] signal: $\Pr(h=k|x)$ for $k=6,7,\dots,N$, wherein the sequence before 5' of the polyadenylation site is $x=(x_1, x_2, \dots, x_N)$ and where x_N is the most 3'-most base before upstream of the polyadenylation site.

10. (original) The method according to Claim 9 wherein $\Pr(h=k|x) = \Pr(x|h=k) \Pr(h=k) / \Pr(x)$.

11. (original) The method of Claim 10 wherein $\Pr(h=k|x) = \Pr(x_{k-5}, \dots, x_k|h=k) \Pr(h=k) / \Pr(x_{k-5}, \dots, x_k)$ and wherein $\Pr(h=k)$ is the probability that the polyadenylation hexamer is located at position k in the sequence, at a distance $(N-k)$ from the polyadenylation site, $\Pr(x_{k-5}, \dots, x_k|h=k)$ is the probability of observing the hexamer (x_{k-5}, \dots, x_k) given that it is a polyadenylation signal and $\Pr(x_{k-5}, \dots, x_k|h \neq k)$ is the probability of observing the hexamer given that it is not from the polyadenylation signal.

12. (currently amended) The method of Claim 11 wherein the step of detecting comprises using a gamma function to produce a density distribution which places the majority of its weight on the positions located 5 to 25 bases distant from the polyadenylation site.

Application No.: 10/028,416

13. (original) The method of Claim 12 wherein $\Pr(x_{k-5}, \dots, x_k | h \neq k)$, the probability of observing the hexamer given that it is not from a polyadenylation signal, is modeled using a second-order Markov model trained on data collected from human 3'UTRs.

14. (original) The method of Claim 13 wherein $\Pr(x_{k-5}, \dots, x_k | h \neq k) = \Pr(x_{k-5}) \Pr(x_{k-4} | x_{k-5}) \Pr(x_{k-3} | x_{k-5}, x_{k-4}) \Pr(x_{k-2} | x_{k-4}, x_{k-3}) \Pr(x_{k-1} | x_{k-3}, x_{k-2}) \Pr(x_k | x_{k-2}, x_{k-1})$ wherein the first term is a zero-order Markovian probability, the second is a first-order Markovian probability and the remaining four terms are second-order Markovian probabilities.

15. (original) The method of Claim 14 wherein, for a k^{th} -order Markov model, the probability of base b following a word w of length k is estimated by the frequency of the concatenated word (wb) divided by the frequency of the word w , where frequencies are computed from the training dataset of 3'UTR sequences.

16. (original) The method of Claim 15 wherein, for the case $k = 0$ (a zero-order Markovian model), the probability of base b is estimated by its frequency in the dataset divided by the size of the dataset.

17. (currently amended) A computer readable medium comprising computer-executable instructions for performing the method comprising:
inputting a plurality of RNA transcript sequences ~~or sequences derived from RNA transcript sequences~~, wherein at least one sequence has its poly A or poly T tract sequence;

Application No.: 10/028,416

searching for a polyadenylation site, wherein the polyadenylation is an adenine rich region at the 3' end of the sequence or a thymine rich region at the ~~beginning~~ 5' end of the sequence;

detecting the presence of polyadenylation signals neighboring the polyadenylation site by ~~scanning~~ analyzing ~~[[the]]~~ EST or RNA sequences or their corresponding genomic DNA sequences.

18. (currently amended) The computer readable medium of Claim 17 wherein the step of searching for a polyadenylation site ~~comprising~~ comprises ~~scanning~~ analyzing the sequences for adenine rich region at the 3' end of the sequence or a thymine rich region at the ~~beginning~~ 5' end of the sequence.

19. (original) The computer readable medium of Claim 18 wherein the adenine rich region comprises adenine in at least 50% of the region and the thymine rich region comprises thymine in at least 50% of the region.

20. (original) The computer readable medium of Claim 19 wherein the adenine rich region comprises adenine in at least 60% of the region and the thymine rich region comprises thymine in at least 60% of the region.

21. (original) The computer readable medium of Claim 20 wherein the adenine rich region comprises adenine in at least 70% of the region and the thymine rich region comprises thymine in at least 70% of the region.

Application No.: 10/028,416

22. (original) The computer readable medium of Claim 21 wherein the adenine rich region comprises adenine in at least 80% of the region and the thymine rich region comprises thymine in at least 80% of the region.

23. (currently amended) The computer readable medium of Claim 17 wherein a heuristic score $n_A / (n_A + 0.5 * (\max(n_R - 20, 0)))$ is used for detecting adenine rich or thymine rich region; wherein n_A is the number of adenines or thymines in the block, and n_R is the number of bases [[after]] downstream of the block of adenines or thymines thymines to the end of the sequence.

24. (currently amended) A computer readable medium comprising computer-executable instructions for performing the method comprising: searching for a polyadenylation signal hexamer in the sequence before 5' of the polyadenylation site.

25. (currently amended) The computer readable medium of Claim 24 wherein the searching comprises evaluating the probability that there is a polyadenylation [[site]] signal: $\Pr(h=k|x)$ for $k=6,7,\dots,N$, wherein the sequence before 5' of the polyadenylation site is $x=(x_1, x_2, \dots, x_N)$ and where x_N is the 3'-most base before upstream of the polyadenylation site.

26. (original) The computer readable medium of Claim 25 wherein: $\Pr(h=k|x) = \Pr(x|h=k) \Pr(h=k) / \Pr(x)$.

Application No.: 10/028,416

27. (original) The computer readable medium of Claim 26 wherein: $\Pr(h=k|x) = \Pr(x_{k-5}, \dots, x_k|h=k) \Pr(h=k) / \Pr(x_{k-5}, \dots, x_k)$ and wherein $\Pr(h=k)$ is the probability that the polyadenylation hexamer is located at position k in the sequence, at a distance $(N-k)$ from the polyadenylation site, $\Pr(x_{k-5}, \dots, x_k|h=k)$ is the probability of observing the hexamer (x_{k-5}, \dots, x_k) given that it is a polyadenylation signal and $\Pr(x_{k-5}, \dots, x_k|h \neq k)$ is the probability of observing the hexamer that it is not from the polyadenylation signal.

28. (currently amended) The computer readable medium of Claim 27 wherein the step of detecting comprises using a gamma function to produce a density distribution which places the majority of its weight on the positions located 5 to 25 bases distant from the polyadenylation site.

29. (original) The computer readable medium of Claim 28 wherein $\Pr(x_{k-5}, \dots, x_k|h \neq k)$, the probability of observing the hexamer given that it is not from a polyadenylation signal, is modeled using a second-order Markov model trained on data collected from human 3'UTRs.

30. (original) The computer readable medium of Claim 29 wherein $\Pr(x_{k-5}, \dots, x_k|h \neq k) = \Pr(x_{k-5}) \Pr(x_{k-4}|x_{k-5}) \Pr(x_{k-3}|x_{k-5}, x_{k-4}) \Pr(x_{k-2}|x_{k-4}, x_{k-3}) \Pr(x_{k-1}|x_{k-3}, x_{k-2}) \Pr(x_k|x_{k-2}, x_{k-1})$ wherein the first term is a zero-order Markovian probability, the second is a first-order Markovian probability and the remaining four terms are second-order Markovian probabilities.

Application No.: 10/028,416

31. (original) The computer readable medium of Claim 30 wherein, for a k^{th} -order Markov model, the probability of base b following a word w of length k is estimated by the frequency of the concatenated word (wb) divided by the frequency of the word w , where frequencies are computed from the training dataset of 3'UTR sequences.

32. (original) The computer readable medium of Claim 31 wherein, for the case $k = 0$ (a zero-order Markovian model), the probability of base b is estimated by its frequency in the dataset divided by the size of the dataset.

33. (currently amended) A system comprising: a processor; and a memory coupled with the processor, the memory storing a plurality of machine instructions that cause the processor to perform logical steps of the method comprising:
inputting a plurality of RNA transcript sequences or Expressed Sequence Tags (EST)
~~sequences derived from RNA transcript sequences~~, wherein at least one sequence has its
poly A or poly T tract sequence;
searching for a polyadenylation site, wherein the polyadenylation is an adenine rich
region at the 3' end of the sequence or a thymine rich region at the ~~beginning~~ 5' end of
the sequence;
detecting the presence of polyadenylation signals neighboring the polyadenylation site by
~~scanning~~ analyzing ~~[[the]]~~ EST or RNA sequences or their corresponding genomic DNA
sequences.

Application No.: 10/028,416

34. (currently amended) The system of Claim 33 wherein the step of searching for a polyadenylation site ~~comprising~~ comprises scanning analyzing the sequences for adenine rich region at the 3' end of the sequence or a thymine rich region at the ~~beginning~~ 5' end of the sequence.

35. (original) The system of Claim 34 wherein the adenine rich region comprises adenine in at least 50% of the region and the thymine rich region comprises thymine in at least 50% of the region.

36. (original) The system of Claim 35 wherein the adenine rich region comprises adenine in at least 60% of the region and the thymine rich region comprises thymine in at least 60% of the region.

37. (original) The system of Claim 36 wherein the adenine rich region comprises adenine in at least 70% of the region and the thymine rich region comprises thymine in at least 70% of the region.

38. (original) The system of Claim 37 wherein the adenine rich region comprises adenine in at least 80% of the region and the thymine rich region comprises thymine in at least 80% of the region.

39. (currently amended) The system of Claim 33 wherein a heuristic score $n_A / (n_A + 0.5 * (\max(n_R - 20, 0)))$ is used for detecting adenine rich or thymine rich region; wherein:

Application No.: 10/028,416

n_A is the number of adenines or thymines in the block, and n_R is the number of bases
[[after]] downstream of the block of adenines or thymines to the end of the sequence.

40. (currently amended) A system comprising a processor; and a memory coupled with the processor, the memory storing a plurality of machine instructions that cause the processor to perform logical steps of the method for detecting polyadenylation signal in a sequence with a polyadenylation site comprising: searching for a polyadenylation signal hexamer in the sequence before 5' of the polyadenylation site.

41. (currently amended) The system of Claim 40 wherein the searching comprises evaluating the probability that there is a polyadenylation ~~[[site]]~~ signal: $\Pr(h=k|x)$ for $k=6,7,\dots,N$, wherein the sequence before 5' of the polyadenylation site is $x=(x_1, x_2, \dots, x_N)$ and where x_N is the most 3'-most base before upstream of the polyadenylation site.

42. (original) The system of Claim 41 wherein: $\Pr(h=k|x) = \Pr(x|h=k) \Pr(h=k)/\Pr(x)$.

43. (original) The system of Claim 42 wherein $\Pr(h=k|x) = \Pr(x_{k-5}, \dots, x_k|h=k) \Pr(h=k)/\Pr(x_{k-5}, \dots, x_k)$ and wherein $\Pr(h=k)$ is the probability that the polyadenylation hexamer is located at position k in the sequence, at a distance $(N-k)$ from the polyadenylation site, $\Pr(x_{k-5}, \dots, x_k|h=k)$ is the probability of observing the hexamer (x_{k-5}, \dots, x_k) given that it is a polyadenylation signal and $\Pr(x_{k-5}, \dots, x_k|h \neq k)$ is the probability of observing the hexamer given that it is not from the polyadenylation signal.

Application No.: 10/028,416

44. (currently amended) The system of Claim 43 wherein the step of detecting comprises using a gamma function to produce a density distribution which places the majority of its weight on the positions located 5 to 25 bases distant from the polyadenylation site.

45. (original) The system of Claim 44 wherein $\Pr(x_{k-5}, \dots, x_k | h \neq k)$, the probability of observing the hexamer given that it is not from a polyadenylation signal, is modeled using a second-order Markov model trained on data collected from human 3'UTRs.

46. (original) The system of Claim 45 wherein $\Pr(x_{k-5}, \dots, x_k | h \neq k) = \Pr(x_{k-5}) \Pr(x_{k-4} | x_{k-5}) \Pr(x_{k-3} | x_{k-5}, x_{k-4}) \Pr(x_{k-2} | x_{k-4}, x_{k-3}) \Pr(x_{k-1} | x_{k-3}, x_{k-2}) \Pr(x_k | x_{k-2}, x_{k-1})$, wherein the first term is a zero-order Markovian probability, the second is a first-order Markovian probability and the remaining four terms are second-order Markovian probabilities.

47. (original) The system of Claim 46 wherein, for a k^{th} -order Markov model, the probability of base b following a word w of length k is estimated by the frequency of the concatenated word (wb) divided by the frequency of the word w , where frequencies are computed from the training dataset of 3'UTR sequences.

48. (original) The system of Claim 47 wherein, for the case $k = 0$ (a zero-order Markovian model), the probability of base b is estimated by its frequency in the dataset divided by the size of the dataset.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.